

JOURNAL OF NUMBER THEORY 8, 99–108 (1976)

## A Periodicity Lemma in Linear Diophantine Analysis

HANS G. WEIDNER

*Mathematisches Institut der Universität Erlangen–Nürnberg, D-852 Erlangen,  
Bismarckstrasse 1½**Communicated by H. W. Leopoldt*

Received January 23, 1973

Let  $g = (g_1, \dots, g_r) \geq 0$  and  $h = (h_1, \dots, h_r) \geq 0$ ,  $g_\rho, h_\rho \in \mathbb{J}$ , be two vectors of nonnegative integers and let  $\lambda \in \mathbb{J}$ ,  $\lambda \geq 0$ ,  $\lambda \equiv 0 \pmod{d}$ , where  $d$  denotes g.c.d.  $(g_1, \dots, g_r)$ . Define

$$A(\lambda) = A(\lambda; g, h) = \min \left\{ \sum_{\rho=1}^r x_\rho h_\rho : x_\rho \geq 0, x_\rho \in \mathbb{J}, \sum_{\rho=1}^r x_\rho g_\rho = \lambda \right\}.$$

It is shown in this paper that  $A(\lambda)$  is periodic in  $\lambda$  with constant jump. If  $i \in \{1, \dots, r\}$  is such that

$$\det \begin{pmatrix} g_i & h_i \\ g_\rho & h_\rho \end{pmatrix} \geq 0 \quad (\rho = 1, \dots, r),$$

then

$$A(\lambda + g_i) = A(\lambda) + h_i$$

holds true for all sufficiently large  $\lambda$ ,  $\lambda \equiv 0 \pmod{d}$ .

## 1. INTRODUCTION

Let  $r \geq 1$  and let  $g = (g_1, \dots, g_r) \geq 0$ ,  $h = (h_1, \dots, h_r) \geq 0$  be vectors of nonnegative integers:  $g_\rho, h_\rho \in \mathbb{J}$ ;  $g_\rho \geq 0$ ,  $h_\rho \geq 0$  ( $\rho = 1, \dots, r$ ). It is well known (cf. [3]) that the affine hyperplane in  $\mathbb{R}^r$

$$g_1 x_1 + \dots + g_r x_r - \lambda = 0 \tag{1}$$

contains vectors  $(x_1, \dots, x_r)$  with nonnegative integer components  $x_\rho$  provided the “parameter”  $\lambda \in \mathbb{J}$ ,  $\lambda \geq 0$ , is sufficiently large and satisfies  $\lambda \equiv 0 \pmod{d}$ , where  $d$  denotes g.c.d.  $(g_1, \dots, g_r)$ .

Consider the linear function

$$f(x_1, \dots, x_r) = h_1 x_1 + \dots + h_r x_r.$$

It is in some respects useful (cf. [4]) to have information about the minimum of  $f$  on the "nonnegative integer lattice" of the hyperplane (1). Let us denote this minimum for fixed  $g$  and  $h$  by  $A(\lambda)$ .

It is shown in this paper that  $A(\lambda)$  enjoys a "periodicity" property as  $\lambda$  is running through the multiples of  $d$ . If  $i \in \{1, \dots, r\}$  is such that

$$g_i h_\rho - g_\rho h_i \geq 0$$

for  $\rho = 1, \dots, r$  (such an  $i$  exists, as is easily seen), then

$$A(\lambda + g_i) = A(\lambda) + h_i \quad (2)$$

holds for all sufficiently large  $\lambda \equiv 0 \pmod{d}$ .

Hence a complete insight in the values of  $A(\lambda)$  is possible, once the finitely many quantities  $A(\lambda_0 + \tau d)$  ( $\tau = 0, 1, \dots, (g_i/d) - 1$ ) are known for some  $\lambda_0$  large enough. A lower bound  $N$  such that (2) holds for  $\lambda \geq N$  is also given in our main theorem.

## 2. PRELIMINARIES

Let  $r \geq 1$  and let  $g = (g_1, \dots, g_r)$  be a vector with nonnegative integer components; we assume  $g_1 \neq 0$ , which is no restriction for our results.

Define

$$d_1 = g_1, \quad d_\rho = \text{g.c.d.}(g_1, \dots, g_\rho); \quad \hat{d}_{\rho-1} = d_{\rho-1}/d_\rho \quad (\rho = 2, \dots, r), \quad (3)$$

$$N_1 = 0, \quad N_\rho = N_\rho(g_1, \dots, g_\rho) = N_{\rho-1} + (\hat{d}_{\rho-1} - 1)(g_\rho - d_\rho) \quad (4) \\ (\rho = 2, \dots, r).$$

Note that

$$d_{\rho-1} \equiv 0 \pmod{d_\rho}, \quad N_\rho \equiv 0 \pmod{d_\rho} \quad (\rho = 2, \dots, r). \quad (5)$$

LEMMA 1. Let  $\lambda \in \mathbb{J}$  be such that  $\lambda \equiv 0 \pmod{d_r}$  and  $\lambda \geq N_r$ . Then

(a) there exists  $\kappa \in \mathbb{J}$ ,  $0 \leq \kappa \leq \hat{d}_{r-1} - 1$ , such that

$$\lambda - \kappa g_r \equiv 0 \pmod{d_{r-1}} \quad \text{and} \quad \lambda - \kappa g_r \geq N_{r-1} \quad (r \geq 2);$$

(b)  $\lambda$  admits a "nonnegative  $g$ -representation"  $(a_1, \dots, a_r)$ , i.e.,  $a_\rho \in \mathbb{J}$  ( $\rho = 1, \dots, r$ ) satisfy

$$a_\rho \geq 0, \quad \sum_{\rho=1}^r a_\rho g_\rho = \lambda.$$

*Proof* (cf. [3]). Put  $\hat{g}_r = g_r/d_r \in \mathbb{J}$ ; then the numbers

$$0, \hat{g}_r, 2\hat{g}_r, \dots, (\hat{d}_{r-1} - 1)\hat{g}_r$$

span all residue classes mod  $\hat{d}_{r-1}$  since any relation

$$p\hat{g}_r \equiv q\hat{g}_r \pmod{\hat{d}_{r-1}}$$

with  $p, q \in \mathbb{J}$  satisfying w.l.o.g.  $0 \leq q < p \leq \hat{d}_{r-1} - 1$ , would imply  $(p - q)\hat{g}_r = l\hat{d}_{r-1}$  for some  $l \in \mathbb{J}$  and  $0 < p - q < \hat{d}_{r-1}$ , which is impossible, since  $d_r = \text{g.c.d.}(d_{r-1}, g_r)$ , i.e.,  $1 = \text{g.c.d.}(\hat{d}_{r-1}, \hat{g}_r)$ .

The integers

$$\begin{aligned} (\hat{d}_{r-1} - 1)\hat{g}_r, (\hat{d}_{r-1} - 1)\hat{g}_r - 1, \dots, (\hat{d}_{r-1} - 1)\hat{g}_r - (\hat{d}_{r-1} - 1) \\ = (\hat{d}_{r-1} - 1)(\hat{g}_r - 1) \end{aligned}$$

also span all residue classes mod  $\hat{d}_{r-1}$ . Therefore for any  $\lambda' \geq (\hat{d}_{r-1} - 1)(\hat{g}_r - 1)$  there is  $\kappa \in \mathbb{J}$ ,  $0 \leq \kappa \leq \hat{d}_{r-1} - 1$  such that

$$\lambda' - \kappa\hat{g}_r \equiv 0 \pmod{\hat{d}_{r-1}}, \quad \lambda' - \kappa\hat{g}_r \geq 0. \quad (6)$$

Let  $\lambda' = (\lambda - N_{r-1})/d_r$ ;  $\lambda'$  is an integer in view of (5) and satisfies  $\lambda' \geq (\hat{d}_{r-1} - 1)(\hat{g}_r - 1)$  by (4). Hence there is  $\kappa$  as in (6), i.e.,

$$\lambda - N_{r-1} - \kappa g_r \equiv 0 \pmod{d_{r-1}}, \quad \lambda - N_{r-1} - \kappa g_r \geq 0.$$

This proves part (a) of the lemma; (b) follows immediately from (a) by an inductive argument.

LEMMA 2. Let  $r \geq 2$ ,  $g = (g_1, \dots, g_r) \geq 0$ ,  $g_o \in \mathbb{J}$ ,  $g_1 \neq 0$ ; let  $\lambda \equiv 0 \pmod{d_r}$ ,  $\lambda \geq N_r$ , and let  $(0, a_2, \dots, a_r)$  be a nonnegative  $g$ -representation of  $\lambda + g_1$ :

$$\lambda + g_1 = a_2 g_2 + \dots + a_r g_r. \quad (7)$$

Then there is a nonnegative  $g$ -representation  $(c_1, \dots, c_r)$  of  $\lambda$

$$\lambda = c_1 g_1 + c_2 g_2 + \dots + c_r g_r \quad (8)$$

such that

$$\sum_{o=i+1}^r (a_o - c_o) g_o \geq 0 \quad (i = 1, \dots, r-1). \quad (9)$$

*Proof.* We proceed by induction on  $r$ .

Let  $r = 2$ ,  $\lambda + g_1 = a_2 g_2$ , and let  $\lambda = c_1 g_1 + c_2 g_2$  be any nonnegative  $g$ -representation of  $\lambda$  that exists by Lemma 1 since  $\lambda \geq N_2$ . Then

$$(a_2 - c_2) g_2 = \lambda + g_1 - \lambda + c_1 g_1 = (1 + c_1) g_1 > 0.$$

Assume now that  $r \geq 3$  and that the lemma is true for all  $j$ ,  $2 \leq j \leq r - 1$ . Let  $\kappa = (\kappa_1, \dots, \kappa_r)$  be the  $g$ -representation of  $\lambda$  obtained by applying Lemma 1 (a)  $(r - 1)$ -times. Then

$$\lambda - \sum_{\rho=i+1}^r \kappa_\rho g_\rho \geq N_i \quad (i = 1, \dots, r - 1) \quad (10)$$

holds true by the very construction of  $\kappa$ . If  $\kappa$  satisfies

$$\sum_{\rho=i+1}^r (a_\rho - \kappa_\rho) g_\rho \geq 0 \quad (i = 1, \dots, r - 1), \quad (11)$$

we set  $c_\rho = \kappa_\rho$  ( $\rho = 1, \dots, r$ ) and are done. If (11) is false, there is  $j$ ,  $2 \leq j \leq r - 1$ , such that

$$\sum_{\rho=j+1}^r (a_\rho - \kappa_\rho) g_\rho < 0. \quad (12)$$

(Note that  $j = 1$  is impossible since

$$\sum_{\rho=2}^r (a_\rho - \kappa_\rho) g_\rho = \lambda + g_1 - (\lambda - \kappa_1 g_1) = (1 + \kappa_1) g_1 > 0.)$$

Define

$$\tilde{\lambda} = \sum_{\rho=2}^j a_\rho g_\rho - g_1.$$

Then  $\tilde{\lambda} \equiv 0 \pmod{d_j}$ ; moreover, by (7), (10), and (12),

$$\begin{aligned} \tilde{\lambda} &= \lambda - \sum_{\rho=j+1}^r a_\rho g_\rho \\ &= \lambda - \sum_{\rho=j+1}^r \kappa_\rho g_\rho - \sum_{\rho=j+1}^r (a_\rho - \kappa_\rho) g_\rho \\ &> N_j. \end{aligned} \quad (13)$$

Since

$$\tilde{\lambda} + g_1 = a_2 g_2 + \dots + a_j g_j,$$

we are in a position to apply the induction hypothesis, thus obtaining a nonnegative vector  $(\tilde{c}_1, \dots, \tilde{c}_j)$  such that

$$\tilde{\lambda} = \sum_{\rho=1}^j \tilde{c}_\rho g_\rho, \quad (14)$$

$$\sum_{\rho=i+1}^j (a_\rho - \tilde{c}_\rho) g_\rho \geq 0 \quad (i = 1, \dots, j-1). \quad (15)$$

Then (13) and (14) provide a representation

$$\lambda = \sum_{\rho=1}^j \tilde{c}_\rho g_\rho + \sum_{\rho=j+1}^r a_\rho g_\rho.$$

Put

$$\begin{aligned} c_\rho &= \tilde{c}_\rho, & \rho &= 1, \dots, j, \\ &= a_\rho, & \rho &= j+1, \dots, r; \end{aligned}$$

then (8) is satisfied, and (9) follows immediately from (15). This completes the proof.

The following simple lemma is stated without proof.

LEMMA 3. Let  $r \geq 2$  and let for  $g = (g_1, \dots, g_r)$ ,  $h = (h_1, \dots, h_r)$

$$D_{\rho, \sigma} = g_\rho h_\sigma - g_\sigma h_\rho = \det \begin{pmatrix} g_\rho & h_\rho \\ g_\sigma & h_\sigma \end{pmatrix} \quad (\rho, \sigma = 1, \dots, r).$$

Then

(a) there is  $i \in \{1, \dots, r\}$  satisfying

$$D_{i, \rho} \geq 0 \quad (\rho = 1, \dots, r),$$

(b) there is an ordering  $\{i_1, \dots, i_r\}$  of  $\{1, \dots, r\}$  such that

$$D_{i_\rho, i_{\rho+1}} \geq 0 \quad (\rho = 1, \dots, r-1).$$

### 3. A PERIODICITY THEOREM

Let  $g = (g_1, \dots, g_r) \geq 0$ ,  $g_\rho \in \mathbb{J}$ , and let  $\lambda \equiv 0 \pmod{d_r}$ . We introduce a short notation for the set of nonnegative  $g$ -representations of  $\lambda$ :

$$\mathfrak{R}(\lambda; g) = \left\{ a = (a_1, \dots, a_r) : \sum_{\rho=1}^r a_\rho g_\rho = \lambda, a_\rho \in \mathbb{J}, a_\rho \geq 0, \rho = 1, \dots, r \right\}.$$

Suppose  $\mathfrak{A}(\lambda; g) \neq \emptyset$ . Let  $h = (h_1, \dots, h_r) \geq 0$ ,  $h_\rho \in \mathbb{J}$ , and define

$$\Lambda(\lambda) = \Lambda(\lambda; g, h) = \min_{a \in \mathfrak{A}(\lambda; g)} \sum_{\rho=1}^r a_\rho h_\rho. \quad (16)$$

Let  $D_{\rho, \sigma} = g_\rho h_\sigma - g_\sigma h_\rho$  ( $\rho, \sigma = 1, \dots, r$ ) and observe Lemma 3.

**THEOREM.** Let  $g = (g_1, \dots, g_r) \geq 0$ ,  $h = (h_1, \dots, h_r) \geq 0$ , be such that

$$D_{1, \rho} \geq 0 \quad (\rho = 1, \dots, r)$$

and

$$D_{\rho, \rho+1} \geq 0 \quad (\rho = 1, \dots, r-1). \quad (17)$$

Then, for every  $\lambda \geq N_r(g_1, \dots, g_r) = \sum_{\rho=2}^r (d_{\rho-1} - 1)(g_\rho - d_\rho)$ ,  $\lambda \equiv 0 \pmod{d_r}$ , the identity

$$\Lambda(\lambda + g_1) = \Lambda(\lambda) + h_1 \quad (18)$$

holds true.

*Proof.* It is obvious that the statement of the theorem is not affected if we omit all  $g_i$  with  $g_i = 0$ . Hence we may assume w.l.o.g.  $g_\rho > 0$  ( $\rho = 1, \dots, r$ ) in the following proofs. Since the theorem is trivial for  $r = 1$ , we may also assume  $r \geq 2$ .

(a) Since  $\lambda \geq N_r$ , we have  $\mathfrak{A}(\lambda; g) \neq \emptyset$  by Lemma 1. Let

$$\mathfrak{A}_0(\lambda; g) = \{a = (a_1, \dots, a_r) \in \mathfrak{A}(\lambda; g) : a_1 = 0\}$$

$$\mathfrak{A}_1(\lambda; g) = \{a = (a_1, \dots, a_r) \in \mathfrak{A}(\lambda; g) : a_1 \geq 1\}.$$

Obviously

$$\mathfrak{A}(\lambda; g) = \mathfrak{A}_0(\lambda; g) \cup \mathfrak{A}_1(\lambda; g). \quad (19)$$

Suppose  $\mathfrak{A}_0(\lambda + g_1; g) \neq \emptyset$  (otherwise the parts (a)–(c) of this proof are dispensable), and let  $a \in \mathfrak{A}_0(\lambda + g_1; g)$ , i.e.,

$$\lambda + g_1 = \sum_{\rho=2}^r a_\rho g_\rho. \quad (20)$$

We shall now prove that

$$\sum_{\rho=2}^r a_\rho h_\rho \geq \Lambda(\lambda) + h_1. \quad (21)$$

To this end let  $c = (c_1, \dots, c_r) \in \mathfrak{A}(\lambda; g)$  be such that

$$\sum_{\rho=i+1}^r (a_\rho - c_\rho) g_\rho \geq 0 \quad (i = 1, \dots, r-1) \quad (22)$$

in accordance with Lemma 2.

(b) We claim

$$D_{1,i} \sum_{\rho=i}^r (a_\rho - c_\rho) g_\rho \geq g_i \sum_{\rho=1}^{i-1} (c_\rho - a_\rho) D_{1,\rho} \quad (i = 2, \dots, r). \quad (23)$$

Before proving this we note that

$$g_i D_{j,k} + g_j D_{k,i} + g_k D_{i,j} = 0, \quad (i, j, k \in \{1, \dots, r\});$$

in particular,

$$g_i D_{1,i+1} + g_1 D_{i+1,i} + g_{i+1} D_{i,1} = 0 \quad (i = 1, \dots, r-1),$$

which implies

$$g_i D_{1,i+1} \geq g_{i+1} D_{1,i} \quad (i = 1, \dots, r-1) \quad (24)$$

in view of (17).

Inequality (23) is shown by induction on  $i$ . If  $i = 2$ , we have

$$\begin{aligned} D_{1,2} \sum_{\rho=2}^r (a_\rho - c_\rho) g_\rho &= D_{1,2}(\lambda + g_1 - (\lambda - c_1 g_1)) \\ &= D_{1,2}(1 + c_1) g_1 \geq 0 = g_2(c_1 - a_1) D_{1,1}. \end{aligned}$$

Suppose now that  $i \leq r-1$  and that (23) holds. Then, by (22), (24), and the induction hypothesis,

$$\begin{aligned} &D_{1,i+1} \sum_{\rho=i+1}^r (a_\rho - c_\rho) g_\rho \\ &\geq (g_{i+1}/g_i) D_{1,i} \sum_{\rho=i+1}^r (a_\rho - c_\rho) g_\rho \\ &= (g_{i+1}/g_i) D_{1,i} \sum_{\rho=i}^r (a_\rho - c_\rho) g_\rho - g_{i+1} D_{1,i}(a_i - c_i) \\ &\geq (g_{i+1}/g_i) g_i \sum_{\rho=1}^{i-1} (c_\rho - a_\rho) D_{1,\rho} - g_{i+1} D_{1,i}(a_i - c_i) \\ &= g_{i+1} \sum_{\rho=1}^i (c_\rho - a_\rho) D_{1,\rho}. \end{aligned}$$

Note that (23) yields for  $i = r$  (recall that  $D_{1,1} = 0$ )

$$\sum_{\rho=2}^r (a_\rho - c_\rho) D_{1,\rho} \geq 0. \quad (25)$$

(c) Observing that  $\sum_{\rho=1}^r c_\rho g_\rho = \lambda$ , we have

$$\begin{aligned} \sum_{\rho=1}^r c_\rho h_\rho &= c_1 h_1 + \sum_{\rho=2}^r c_\rho h_\rho = (1/g_1) \left( \lambda - \sum_{\rho=2}^r c_\rho g_\rho \right) h_1 + \sum_{\rho=2}^r c_\rho h_\rho \\ &= (1/g_1) \left( \lambda h_1 + \sum_{\rho=2}^r c_\rho D_{1,\rho} \right). \end{aligned}$$

Similarly, by (20),

$$\sum_{\rho=2}^r a_\rho h_\rho = (1/g_1) \left( (\lambda + g_1) h_1 + \sum_{\rho=2}^r a_\rho D_{1,\rho} \right).$$

Hence, by (25),

$$\sum_{\rho=2}^r a_\rho h_\rho - \sum_{\rho=1}^r c_\rho h_\rho = (1/g_1) \left( g_1 h_1 + \sum_{\rho=2}^r (a_\rho - c_\rho) D_{1,\rho} \right) \geq h_1,$$

which provides (21):

$$\sum_{\rho=2}^r a_\rho h_\rho \geq \sum_{\rho=1}^r c_\rho h_\rho + h_1 \geq \Lambda(\lambda) + h_1.$$

(d) Now let  $a = (a_1, \dots, a_r) \in \mathfrak{A}_1(\lambda + g_1; g)$ ; then  $(a_1 - 1, a_2, \dots, a_r) \in \mathfrak{A}(\lambda; g)$ , and we obtain at once

$$\sum_{\rho=1}^r a_\rho h_\rho = (a_1 - 1) h_1 + \sum_{\rho=2}^r a_\rho h_\rho + h_1 \geq \Lambda(\lambda) + h_1. \quad (26)$$

(e) In view of (19) the inequalities (21) and (26) show that

$$\Lambda(\lambda + g_1) \geq \Lambda(\lambda) + h_1.$$

On the other hand, let  $b = (b_1, \dots, b_r) \in \mathfrak{A}(\lambda; g)$  represent  $\Lambda(\lambda)$ , i.e.,  $\sum_{\rho=1}^r b_\rho h_\rho = \Lambda(\lambda)$ . Then obviously  $(b_1 + 1, b_2, \dots, b_r) \in \mathfrak{A}(\lambda + g_1; g)$ , which means

$$\Lambda(\lambda + g_1) \leq (b_1 + 1) h_1 + \sum_{\rho=2}^r b_\rho h_\rho = \Lambda(\lambda) + h_1.$$

This completes the proof.



Next we prove a lemma in order to add a generalizing corollary to the theorem.

LEMMA 4. *Let  $p > 0$ ,  $p \equiv 0 \pmod{d_r}$  be minimal in the sense that there exist integers  $N \geq 0$  and  $q \geq 0$  satisfying*

$$\Lambda(\lambda + p) = \Lambda(\lambda) + q \quad (27)$$

*for every  $\lambda \equiv 0 \pmod{d_r}$ ,  $\lambda \geq N$ . If  $p' > 0$ ,  $p' \equiv 0 \pmod{d_r}$  is such that*

$$\Lambda(\lambda + p') = \Lambda(\lambda) + q' \quad (28)$$

*holds true for some  $q' \geq 0$  and every  $\lambda \equiv 0 \pmod{d_r}$ ,  $\lambda \geq N'$ , then there is  $k \in \mathbb{J}$  such that*

$$p' = kp, \quad q' = kq.$$

*Proof.* Let  $k > 0$  and  $\hat{p}$ ,  $0 \leq \hat{p} < p$ , be such that  $p' = kp + \hat{p}$ , where  $\hat{p} \equiv 0 \pmod{d_r}$ . Let  $\lambda \geq \tilde{N} = \max(N - \hat{p}, N')$ ,  $\lambda \equiv 0 \pmod{d_r}$ . Then

$$\Lambda(\lambda + p') = \Lambda(\lambda + \hat{p} + kp) = \Lambda(\lambda + \hat{p}) + kq,$$

which, in view of (28), means that

$$\Lambda(\lambda + \hat{p}) = \Lambda(\lambda) + q' - kq, \quad (29)$$

where  $\hat{q} = q' - kq \geq 0$ , as is easily seen. Equation (29) contradicts the minimality property of  $p$  except for  $\hat{p} = 0$ , which implies  $\hat{q} = 0$ .

COROLLARY. *Let  $g$  and  $h$  be as in the Theorem and assume*

$$D_{1,\rho} = 0 \quad (\rho = 1, \dots, j) \quad (30)$$

*for some  $j$ ,  $1 \leq j \leq r$ . Then*

$$\Lambda(\lambda + d_j) = \Lambda(\lambda) + (h_1/g_1) d_j$$

*holds for sufficiently large  $\lambda$ ,  $\lambda \equiv 0 \pmod{d_r}$ .*

*Proof.* The corollary follows immediately from the theorem and from Lemma 4, since (30) implies

$$D_{i,\rho} \geq 0 \quad (\rho = 1, \dots, j)$$

for every  $i \in \{1, \dots, j\}$ .

We remark that  $(h_1/g_1) d_j \in \mathbb{J}$ . We have, by (30),

$$\frac{g_1}{d_j} h_p = \frac{g_p}{d_j} h_1, \quad \text{i.e.,} \quad \frac{g_1}{d_j} \mid \frac{g_p}{d_j} h_1 \quad (p = 1, \dots, j);$$

hence  $g_1/d_j$  is a divisor of  $\text{g.c.d.}(g_1 h_1/d_j, \dots, g_j h_1/d_j) = h_1$ .

#### REFERENCES

1. R. D. CARMICHAEL, "The Theory of Numbers and Diophantine Analysis," Dover, New York, 1959.
2. G. H. HARDY AND E. M. WRIGHT, "An Introduction to the Theory of Numbers," Clarendon Press, Oxford, 1954.
3. J. ROSENMÜLLER AND H. G. WEIDNER, A class of extreme convex set functions with finite carrier, *Advances in Math.* **10** (1973), 1-38.
4. J. ROSENMÜLLER AND H. G. WEIDNER, Extreme convex set functions with finite carrier: General theory, *Discrete Mathematics* **10** (1974), 343-382.